

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Samara, in 12 villages there have occurred 64 cases of cholera. In the city of Zarizyn there have been registered 21 cholera cases (and 17 deaths).

In the Trans-Caspian district there were registered during the period from November 8 to 16, 3 fresh cases of plague in Serachs.

According to supplementary official reports, there have been registered in Baku between August 28 and November 13 a total number of 385 cases of cholera (with 211 deaths).

Turkey.—According to Official Bulletin No. 45 of November 21, there have been registered 84 new cases of cholera (and 63 deaths).

Death rate of Berlin and other cities.

The death rate of Berlin for the week ended November 19 was higher than it has been since the middle of September, amounting, calculated on the year, to 14.4 per 1,000 of the population, and thus corresponded almost exactly with the rate for the same week of last year. two-thirds of the large towns and cities of Germany showed more unfavorable health conditions than Berlin, the rate of mortality in the following places being considerably higher than that of this city, viz: Hamburg, Cologne, Düsseldorf, Magdeburg, Halle, Leipzig, Munich, Breslau, Danzig, Königsberg, as well as London, Paris, and Vienna. On the other hand, the death rate of the following cities was lower than the Berlin figure, namely: Dresden, Bremen, Frankfort-on-the-Main, Carlsruhe, Rixdorf (with 13.2), Charlottenburg (11.1), and Schöneberg (8.7). The number of deaths among children in the first year again showed a decrease as compared with the preceding week, so that the increased number of deaths occurred exclusively among the higher-age classes. The death rate of infants, 3.2 per year and 1,000, was lower than the Hamburg, Munich, and Leipzig figures. Acute diseases of the respiratory organs showed a decrease this week, causing 48 deaths (including 3 from influenza). Acute intestinal diseases showed no important variation, claiming 36 victims, while 8 persons succumbed to cholerine. Furthermore, there were registered 84 deaths from phthisis pulmonalis, 40 deaths from cancer, 11 deaths from scarlet fever, 8 deaths from diphtheria, 4 deaths from measles, and 2 deaths from enteric fever. Finally, 3 persons died by violence.

INDIA.

Reports from Bombay—Cholera, plague, and smallpox mortality— Transactions of inspection service.

Acting Asst. Surg. Edward H. Hume reports, November 26, as follows:

Mortality of Bombay city.

| | Nov. 8. | Nov. 15. | Nov. 22. |
|--|-----------------------------|--|--|
| Plague attacks Plague deaths Plague mortality per 1,000 Smallpox deaths Cholera deaths | 52 3.48 2 0 609 | b 82 75 5.02 7 1 638 42.75 | 6 70 57 3.81 2 0 635 42.55 |
| Total deaths. Total mortality per 1,000 | | | |